

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
17 January 2002 (17.01.2002)

PCT

(10) International Publication Number
WO 02/04433 A2

- (51) International Patent Classification⁷: C07D 295/096, A61K 31/495, A61P 3/04
- (74) Agent: DOCTER, Stephen, H.; McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker, Suite 3200, Chicago, IL 60606 (US).
- (21) International Application Number: PCT/US01/41289
- (22) International Filing Date: 6 July 2001 (06.07.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/216,081 6 July 2000 (06.07.2000) US
- (71) Applicant (for all designated States except US): NEUROGEN CORPORATION [US/US]; 35 Northeast Industrial Road, Branford, CT 06405 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): BAKTHAVATCHALAM, Rajagopal [US/US]; 92 Village Lane, Branford, CT 06405 (US). HUTCHISON, Alan [US/US]; 29 Kimberly Lane, Madison, CT 06443 (US). THURKAUF, Andrew [US/US]; 16 Fox Den Road, Danbury, CT 06811 (US).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MELANIN CONCENTRATING HORMONE RECEPTOR LIGANDS



(57) Abstract: Disclosed are compounds of formula (I), and the pharmaceutically acceptable salts thereof wherein Q, X, Y, Z, and R₁ R₉, and R₁₂-R₁₉ are defined herein. These compounds are selective modulators of MCH 1 receptors that are, therefore, useful in the treatment of a variety of metabolic, feeding, and sexual disorders. Methods of treatment of such disorders and well as packaged pharmaceutical compositions are also disclosed.

WO 02/04433 A2

Melanin Concentrating Hormone Receptor Ligands

Background of the Invention

5 This application claims priority from U.S. Provisional Application S.N. 60/216,081, filed July 6, 2000.

Field of the invention

 This invention relates to phenylcycloalkylmethylamino and phenylalkenylamino derivatives, including 1-phenyl-2-aminomethylcyclopropanes, that are modulators of melanin
10 concentrating hormone type 1 (MCH 1) receptors. This invention also relates to pharmaceutical compositions comprising such compounds.

Description of the Related Art

 Melanin concentrating hormone, or MCH, is a cyclic 19 amino acid neuropeptide that
15 is produced within the hypothalamus of many vertebrate species including man. I.C.V. injection of MCH into the lateral ventricle of the hypothalamus has been shown to increase caloric consumption in rats over similarly treated control animals. Furthermore, rats having the *ob/ob* genotype exhibit a 50-80% increase in MCH mRNA expression as compared to leaner *ob/+* genotype mice. MCH knockout mice are leaner than their MCH-producing
20 siblings due to hypophagia and an increased metabolic rate. Thus, MCH is thought to be an important regulator of feeding behavior and body weight.

 The MCH 1 receptor was originally obtained from human cDNA and genomic libraries and characterized as a 402 amino acid G-coupled protein receptor having substantial sequence identity to the somatostatin receptors. This receptor was named the SLC-1 receptor.
25 A rat orthologue of the MCH 1 receptor was isolated from a rat brain cDNA library by Lakaye, et al. (BBA (1998) 1401: 216-220) and found to encode a 353 amino acid protein having seven transmembrane alpha helices and three consensus *N*-glycosylation sites. The rat MCH 1 receptor reported by Lakaye also disclosed was homologous to the human MCH 1 receptor disclosed earlier except for the removal of a 5' intron. Accordingly, Lakaye, et al.,
30 deduced the "corrected" amino acid sequence of the N-terminus of MCH 1 receptor is found within a sequence deposited for a 128 kb fragment of human chromosome 22 encompassing the earlier disclosed MCH 1 receptor gene (Genbank accession number: Z86090).

The earlier reported 402 amino acid MCH 1 receptor protein does not interact with MCH. Thus, the 353 amino acid receptor first reported by Lakaye, is now considered to be the correct full-length sequence for the human MCH 1 receptor.

Immunohistochemistry studies of rat brain sections indicate that the MCH 1 receptor is
5 widely expressed in the brain. MCH 1 receptor expression was found in the olfactory tubercle, cerebral cortex, substantia nigra, basal forebrain CA1, CA2, and CA3 field of the hippocampus, amygdala, and in nuclei in the hypothalamus, thalamus, midbrain and hindbrain. Strong signals have been observed in the ventromedial and dorsomedial nuclei of the hypothalamus, two areas of the brain known to be involved in feeding behavior.

10 Upon binding MCH, MCH 1 receptors expressed in HEK 293 cell mediate a dose dependent release of intracellular calcium. Cells expressing MCH receptors have also been shown to exhibit a pertussis toxin sensitive dose-dependent inhibition of forskolin-elevated cyclic AMP, indicating that the receptor couples to a $G_{i/o}$ G-protein alpha subunit.

Because MCH has been shown to be an important regulator of food intake and energy
15 balance, ligands capable of modulating the activity of the MCH 1 receptor are highly desirable for the treatment of eating disorders and metabolic disorders. Orally available, small molecule, non-peptide antagonists of the MCH 1 receptor are particularly sought for the treatment of obesity.

20

SUMMARY OF THE INVENTION

The invention provides novel compounds, particularly phenylcycloalkylmethylamino and phenylalkenylamino compounds, including 1-phenyl-2-aminomethylcyclopropanes, that are small molecule MCH receptor ligands, especially MCH 1 receptor ligands, that are non-peptide and amino acid free, which compounds exhibit a K_i at the MCH receptor of less than 1 micromolar. Preferred MCH 1 receptors are mammalian receptors, including human and monkey MCH receptors and may either be cloned, recombinantly expressed receptors or naturally expressed receptors.

10 In certain embodiments these compounds also possess one or more, and preferably two or more, three or more, or all of the following properties in that they are: 1) multi-aryl in structure (having a plurality of un-fused or fused aryl groups), 2) orally available *in vivo* (such that a sub-lethal or pharmaceutically acceptable oral dose can provide a detectable *in vivo* effect such as a reduction of appetite), 3) capable of inhibiting the binding of MCH to the MCH receptor at nanomolar concentrations or 4) capable of inhibiting the binding of MCH to the MCH receptor at sub-nanomolar concentrations.

The invention also provides novel compounds of Formula I, shown below, that bind specifically, and preferably with high affinity, to MCH receptors.

20 The invention also provides pharmaceutical compositions comprising compounds of Formula I together with at least one pharmaceutically acceptable carrier. The compounds are particularly useful in the treatment of metabolic, feeding, and sexual disorders. The invention further comprises a method of treating a patient in need of such treatment with a sufficient concentration of a compound of the invention. A preferred concentration is one sufficient to inhibit the binding of MCH to MCH 1 receptors *in vitro*. Treatment of humans, domesticated companion animals (pets) or livestock animals suffering such conditions with an effective amount of a compound of the invention is contemplated by the invention.


Also included in the invention are methods of treating eating disorders, particularly obesity and bulimia nervosa, comprising administering to a patient in need of such treatment a MCH 1 receptor modulator together with leptin, a leptin receptor agonist, or a melanocortin receptor 4 (MC4) agonist.

30 In a separate aspect, the invention provides methods of using compounds of this invention as positive controls in assays for receptor activity and using appropriately labeled compounds of the invention as probes for the localization of receptors, particularly MCH receptors, in tissue sections.

The invention provides compounds and compositions that are useful as inhibitors of MCH binding to MCH 1 receptor, and as inhibitors of MCH mediated signal transduction (e.g., they may be used as standards in assays of MCH binding and MCH-mediated signal transduction). The invention additionally comprises methods of inhibiting MCH binding to

5 MCH receptors *in vivo*, preferably MCH 1 receptors present in the hypothalamus.


Accordingly, a broad embodiment of the invention is directed to a compounds and pharmaceutically acceptable salts of Formula I:



I

wherein:

Q is a group of the Formula:



wherein

15 A is C₁-C₅ alkylene optionally mono-, di, or trisubstituted with substituents independently chosen from C₁-C₃ alkyl, C₁-C₃ alkoxy, halogen, halo(C₁-C₃)alkyl, halo(C₁-C₃)alkoxy, hydroxy, amino, and mono- or di(C₁-C₃)alkylamino;

R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are the same or different and represent hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -SO₂NH₂, mono or dialkylsulfonamido, -C(O)NH₂, or mono or di(C₁-C₆)alkylcarboxamido;

20

R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ independently represent hydrogen or C₁-C₆ alkyl;

25 W is nitrogen or C-R_a where R_a represents hydrogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl or cyano;

X represents halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkylamino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di(C₁-C₆)alkylcarboxamido, -SO₂NH₂, mono or di(C₁-C₆)alkylsulfonamido; or

- 5 X represents phenyl which may be optionally substituted by up to five substituents, which may be the same or different and are selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di-(C₁-C₆)alkylcarboxamido, -SO₂NH₂, and mono or di(C₁-C₆)alkylsulfonamido;
- 10

Y is oxygen, sulfur, -S(O)-, or -SO₂-; and

Z is C₁-C₆ alkyl or mono, di or trifluoromethyl.

The invention also provides intermediates and methods useful for preparing the compounds of Formula I.


15

DETAILED DESCRIPTION OF THE INVENTION

The invention particularly includes compounds and salts of Formula I wherein Q is a ring and A is methylene optionally substituted with C₁-C₂ alkyl.

5 The invention is also specifically directed to compounds and salts of Formula I wherein W is nitrogen or CH and A is methylene. Preferred compounds and salts of this class are those wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen. Other preferred compounds and salts of this class are those wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen, X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl. Also
 10 preferred are compounds and salts of Formula I wherein W is nitrogen or CH and A is methylene, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen, R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy; X is hydrogen, halogen, or phenyl, or most preferably X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

15 Particularly provided by the invention are compounds of Formula II



II

and the pharmaceutically acceptable salts thereof; wherein

A is methylene optionally substituted with C₁-C₂ alkyl, and R₁-R₁₉, W, X, Y, and Z are as
 20 defined for Formula I.

Preferred compounds and salts of Formula II are those wherein W is nitrogen or CH.


Other preferred compounds and salts of Formula II are those wherein W is nitrogen or CH and R₁₀, R₁₁, R₁₂, R₁₃, R₁₅, R₁₇, R₁₈, and R₁₉ are hydrogen.

Also preferred are compounds and salts of Formula II wherein W is nitrogen or CH,
 25 R₁₀, R₁₁, R₁₂, R₁₃, R₁₅, R₁₇, R₁₈, and R₁₉ are hydrogen, R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ independently represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy; R₁₄ and R₁₆ are the same or different and are either hydrogen or methyl; X is hydrogen, halogen, or phenyl; Y is oxygen; and Z is C₁-C₆ alkyl.

Particularly preferred compounds and salts of Formula II are those wherein W is nitrogen or CH, R₁, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen,

R₂, R₃, R₄, and R₅ are independently hydrogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, or halogen; X is
5 halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

The invention further provides compounds of Formula III



III

10 and the pharmaceutically acceptable salts thereof, wherein R₁-R₁₉, W, X, Y, and Z are as defined for Formula I.

Preferred compounds and salts of Formula III are those wherein R₁₃, R₁₅, R₁₇, R₁₉, are hydrogen; and R₁₀, R₁₁, R₁₂, R₁₄, R₁₆, and R₁₈ independently represent hydrogen or methyl, or
15 more preferably hydrogen.


Also preferred are compounds and salts of Formula III, wherein R₁₀-R₁₉ are hydrogen, and W is N or CH.

More preferred compounds and salts of Formula III are those wherein R₁₀-R₁₉ are hydrogen, W is N or CH; R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ independently represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy; X is hydrogen or halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

Particularly preferred compounds and salts of Formula III are those wherein R₁₀-R₁₉ are hydrogen, W is N or CH, R₁, R₂, R₃, R₄ independently represent hydrogen, halogen, C₁-C₂alkyl, or C₁-C₂ alkoxy; R₅, R₆, R₇, and R₈ are hydrogen; X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

Another embodiment of the invention is directed to compounds and salts of Formula

IV



IV

and the pharmaceutically acceptable salts thereof, wherein R₁-R₁₉, W, X, Y, and Z are as defined for Formula I.

5


Preferred compounds and salts of Formula IV are those wherein R₁₃, R₁₅, R₁₇, R₁₉, are hydrogen; and R₁₀, R₁₁, R₁₂, R₁₄, R₁₆, and R₁₈ independently represent hydrogen or methyl, or more preferably hydrogen.

Also preferred are compounds and salts of Formula IV, wherein R₁₀-R₁₉ are hydrogen, and W is N or CH.

More preferred compounds and salts of Formula IV are those wherein R₁₀-R₁₉ are hydrogen, W is N or CH; R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ independently represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy; X is hydrogen or halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

Particularly preferred compounds and salts of Formula IV are those wherein R₁₀-R₁₉ are hydrogen, W is N or CH, R₁, R₂, R₃, R₄ independently represent hydrogen, halogen, C₁-C₂alkyl, or C₁-C₂ alkoxy; R₅, R₆, R₇, and R₈ are hydrogen; X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl.


The invention also provides compounds of Formula V



V

or a pharmaceutically acceptable salt thereof wherein:

Q is a group of the Formula:



wherein:

A is C₁-C₅ alkylene optionally mono-, di, or trisubstituted with substituents independently chosen from C₁-C₃ alkyl, C₁-C₃ alkoxy, halogen, halo(C₁-C₃)alkyl, halo(C₁-C₃)alkoxy, hydroxy, amino, and mono- or di(C₁-C₃)alkylamino;

R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are the same or different and represent hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -SO₂NH₂, mono or dialkylsulfonamido, -C(O)NH₂, or mono or di(C₁-C₆)alkylcarboxamido;

R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ independently represent hydrogen or C₁-C₆ alkyl;

W is nitrogen or C-R_a where R_a represents hydrogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl or cyano;

X represents halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkylamino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di(C₁-C₆)alkylcarboxamido, -SO₂NH₂, mono or di(C₁-C₆)alkylsulfonamido; or


X represents phenyl which may be optionally substituted by up to five substituents, which may be the same or different and are selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di-(C₁-C₆)alkylcarboxamido, -SO₂NH₂, and mono or di(C₁-C₆)alkylsulfonamido;

Y is oxygen, sulfur, -S(O)-, or -SO₂-; and

Z is C₁-C₆ alkyl or mono, di or trifluoromethyl.

Compounds of Formula V are intermediates, useful in preparing compounds MCH 1 receptor ligands.

Preferred compounds of Formula V are those wherein Q is a group the formula



where A is methylene optionally substituted with C₁-C₂ alkyl or A is a single bond. Such compounds will be referred to as compounds of Formula VA.


5 The invention is particularly directed to compounds of Formula VA wherein W is nitrogen or CH.

More preferred compounds of Formula VA are those wherein W is nitrogen or CH, and R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen.

Other preferred compounds of Formula VA are those wherein W is nitrogen or CH,
 10 R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen, X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

Especially preferred compounds of Formula VA are those wherein W is nitrogen or CH, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen, R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆
 15 alkoxy, trifluoromethyl, or trifluoromethoxy; X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

Particularly preferred compounds of Formula V include those where Q is a group of the formula:



where A is methylene optionally substituted with C₁-C₂ alkyl. These compounds are
 20 hereinafter referred to as compounds of Formula VI-A. Specific compounds of Formula VI-A include those where A is methylene and R₁₀ and R₁₁ are methyl or, preferably, hydrogen.

Specific compounds of VA include those wherein W is nitrogen or CH. Preferred compounds of V and VA include those wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen.

25 Other specific compounds of VA include those wherein: X is halogen; Y is oxygen; and Z is C₁-C₆ alkyl.

Still other specific compounds of VA include those where R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are the same or different and represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;



30 X is halogen;





Y is oxygen; and

Z is C₁-C₆ alkyl.

- 5 Preferably not more than 5, and more preferably not more than 3, cyano or nitro groups are present in compounds of Formula I – Formula VA. Preferably not more than 2 of R₁, R₂, R₃, R₄, R₅ are non-hydrogen substituents. Preferably not more than 5 and more preferably not more than 3 of R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are non-hydrogen substituents.

Representative compounds of Formula I are shown in Table 1.

Table 1	
Compound number	Chemical Structure
1	
2	

3	
4	
5	
6	

In certain situations, the compounds of this invention may contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. These compounds can be, for example, racemates or optically active forms. In these situations, the single enantiomers, i.e., optically active forms, can be obtained by asymmetric synthesis or by resolution of the racemates. Asymmetric synthesis of compounds of the invention may be performed using the methods illustrated in Example 1, below. For

compounds having an alpha-methyl benzyl group (R_3 is methyl, R_4 is hydrogen) the R enantiomer is preferred. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent, or chromatography, using, for example a chiral HPLC column.

5 Representative compounds of the invention, which are encompassed by Formula I, include, but are not limited to the compounds in Table I and their pharmaceutically acceptable acid addition salts. In addition, if the compound of the invention is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt. Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically
10 acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds.

Non-toxic pharmaceutical salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic,
15 citric, tartaric, maleic, hydroiodic, alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

The invention also encompasses the acylated prodrugs of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies which may be
20 employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I.

Where a compound exists in various tautomeric forms, the invention is not limited to any one of the specific tautomers. The invention includes all tautomeric forms of a compound.

This invention relates to compounds that bind with high affinity to the melanin
25 concentrating hormone receptors, including human melanin concentrating hormone receptors. This invention also includes such compounds that bind with high selectivity to the melanin concentrating hormone receptors, including human and monkey melanin concentrating hormone receptors. Without wishing to be bound to any particular theory, it is believed that the interaction of the compounds of Formula I with the melanin concentrating hormone
30 receptor results in the pharmaceutical utility of these compounds.

The invention further comprises methods of treating patients in need of such treatment with an amount of a compound of the invention sufficient to alter the symptoms of a disorder.

The diseases and/ or disorders that can also be treated using compounds and compositions according to the invention include, but are not limited to, eating disorders,

sexual disorders, obesity, bulimia, anorexia, diabetes, heart disease, stroke, anorgasmia, or psychogenic impotence.

The invention also provides pharmaceutical compositions comprising at least one compound of the invention together with at least one pharmaceutically acceptable carrier or excipient. Such pharmaceutical compositions include packaged pharmaceutical compositions for treating disorders responsive to melanin concentrating hormone receptor modulation, e.g., treatment of eating disorders such as obesity or bulimia or treatment of sexual disorders such as anorgasmic or psychogenic impotence. The packaged pharmaceutical compositions include a container holding a therapeutically effective amount of at least one melanin concentrating hormone receptor modulator as described supra and instructions (e.g., labeling) indicating that the contained composition is to be used for treating a disorder responsive to melanin concentrating hormone receptor modulation in the patient.

The invention also pertains to methods of inhibiting the binding of melanin concentrating hormone to melanin concentrating hormone receptors which methods involve contacting a compound of the invention with cells expressing melanin concentrating hormone receptors, wherein the compound is present at a concentration sufficient to inhibit melanin concentrating hormone binding to melanin concentrating hormone receptors *in vitro*. This method includes inhibiting the binding of melanin concentrating hormone to melanin concentrating hormone receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to inhibit the binding of melanin concentrating hormone to melanin concentrating hormone receptors *in vitro*. The amount of a compound that would be sufficient to inhibit the binding of melanin concentrating hormone to the melanin concentrating hormone receptor *in vitro* may be readily determined via a melanin concentrating hormone receptor binding assay, such as the assay described in Example 5. The membranes, comprising melanin concentrating hormone receptors, used to determine *in vitro* binding may be obtained from a variety of sources, for example from preparations of HEK 293 cells expressing cloned human or cloned monkey melanin concentrating hormone receptors, especially HEK 293 cells expressing such receptors.

The invention also pertains to methods for altering the signal-transducing activity of MCH receptors, particularly the MCH receptor-mediated release of intracellular calcium, said method comprising exposing cells expressing such receptors to an effective amount of a compound of the invention. This method includes altering the signal-transducing activity of MCH receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to alter the signal-transducing activity of MCH receptors *in vitro*. The

amount of a compound that would be sufficient to alter the signal-transducing activity of MCH receptors may be determined via a MCH receptor signal transduction assay, such as the calcium mobilization assay described in Example 6.

The melanin concentrating hormone receptor ligands provided by this invention and
5 labeled derivatives thereof are also useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the melanin concentrating hormone receptor.

Labeled derivatives the melanin concentrating hormone receptor ligands provided by this invention are also useful as radiotracers for positron emission tomography (PET) imaging or for single photon emission computerized tomography (SPECT).

10 Preferred compounds of the invention do not exhibit fungicidal activity. Such a lack of fungicidal activity may be demonstrated by no more than a 40% reduction of colony size (when treated with the compound at 100 p.p.m. and compared to untreated controls) of *Aspergillus nidulans* strain R153 when grown for 48 hours at 32°C on solid MAG medium. Optionally, BENOMYL, 100 p.p.m., may be used as a positive control. MAG medium is 2%
15 malt extract, 0.2% peptone, 1% glucose and trace elements, pH 6.5. Trace elements as a 5000-fold concentrate consist of 10 g/l EDTA, 4.4 g/l ZnSO₄·7H₂O, 1.01 g/l MnCl₂·4H₂O, 0.32 g/l CoCl₂·6H₂O, 0.315 g/l CuSO₄·5H₂O, 0.22 g/l (NH₄)₆Mo₇O₂₄·H₂O, 1.47 g/l CaCl₂·2H₂O and 1.0 g/l FeSO₄·7H₂O. Medium is made solid by the addition of 1.5% agar.

Alternatively, such a lack of fungicidal activity may be demonstrated by an infection
20 frequency of 60-100% (as compared to untreated plants) for each of *Puccinia recondita* (leaf rust) on wheat, *Erysiphe graminis* (powdery mildew) on barley, *Venturia inaequalis* (scab, black spot) on apple plants, and *Cercospora arachidicola* (early leafspot) on peanut.

The technique employed to determine fungicidal activity is as follows. The plants are grown in John Innes Potting Compost (No. 1, or Seed, as appropriate) in 4 cm diameter mini-
25 pots. A layer of fine sand is placed at the bottom of the pot to facilitate uptake of test compound by the roots.

The test compounds are formulated, e.g., by bead-milling with aqueous Dispersol T or as a solution in acetone/ethanol which is diluted to the required concentration immediately before use. 100 p.p.m. a.i. suspensions are sprayed on to the foilage and applied to the roots of
30 the same plant via the soil. (Sprays are applied to maximum retention, and root drenches to a final concentration equivalent to approximately 40 ppm a.i./dry soil). Tween 20, to give a final concentration of 0.1%, is added when the sprays are applied to the cereals.

For most of the tests, the test compound is applied to the soil (roots) and to the foliage (by spraying) one or two days before the plant is inoculated with the diseases. An

exception is the test on *Erysiphe graminis*, in which the plants are inoculated 24 hours before treatment. After inoculation, the plants are put into an appropriate environment to allow infection to take place and then incubated until the disease is ready for assessment. The period between inoculation and assessment typically varies from 4 to 14 days according to the disease and environment.

Chemical Description and Terminology

The compounds of the invention have asymmetric centers; this invention includes all of the optical isomers and mixtures thereof.

10 Compounds of the invention with carbon-carbon double bonds occur in Z- and E-forms; all isomeric forms of the compounds are included in the invention.

When any variable occurs more than one time in Formula I, its definition on each occurrence is independent of its definition at every other occurrence.

By "C₁-C₆ alkyl" or in the invention is meant straight or branched chain alkyl groups or cycloalkyl groups having 1-6 carbon atoms, such as, for example, methyl, ethyl, propyl, 15 isopropyl, n-butyl, sec-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl, and 3-methylpentyl. Preferred C₁-C₆ alkyl groups are methyl, ethyl, propyl, butyl, cyclopropyl, cyclopropylmethyl, cyclohexyl, cycloheptyl, norbornyl, and the like. Particularly preferred alkyl groups are methyl and ethyl.

20 By "C₁-C₆ alkoxy" in the invention is meant an alkyl group of indicated number of carbon atoms attached through an oxygen bridge such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, 2-pentyl, isopentoxy, neopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy. Preferred alkoxy groups herein are C₁-C₄ alkoxy groups. Particularly preferred alkoxy groups are ethoxy and 25 methoxy.

The term "halogen" includes fluorine, chlorine, bromine, and iodine. Where X is halogen in Formula I- Formula V, bromine is particularly preferred.

"Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more 30 halogen atoms. Examples of haloalkyl include, but are not limited to, mono-, di-, or tri-fluoromethyl, mono-, di-, or tri-chloromethyl, mono-, di-, tri-, tetra-, or penta-fluoroethyl, and mono-, di-, tri-, tetra-, or penta-chloroethyl. Typical haloalkyl groups are trifluoromethyl and

difluoromethyl. Preferably not more than 5, and more preferably not more than 3 haloalkyl groups, are present in compounds of the invention.

"Haloalkoxy" represents a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge.

5 Non-toxic "pharmaceutically acceptable salts" include, but are not limited to salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrite or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, 2-hydroxyethylsulfonate, salicylate and stearate. Similarly, pharmaceutically acceptable cations include, but are not
10 limited to sodium, potassium, calcium, aluminum, lithium and ammonium. The invention also encompasses the prodrugs of the compounds of Formula I.

Pharmaceutical preparations

Those skilled in the art will recognize various synthetic methodologies that may be
15 employed to prepare non-toxic pharmaceutically acceptable prodrugs of the compounds encompassed by Formula I. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable solvents that may be used to prepare solvates of the compounds of the invention, such as water, ethanol, mineral oil, vegetable oil, and dimethylsulfoxide.

The compounds of general Formula I may be administered orally, topically,
20 parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. Oral administration in the form of a pill, capsule, elixir, syrup, lozenge, troche, or the like is particularly preferred. The term parenteral as used herein includes subcutaneous injections, intradermal, intravascular (e.g., intravenous), intramuscular, spinal, intrathecal injection or
25 like injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of
30 general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may

contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These
5 excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and
10 absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium
15 phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose,
20 sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters
25 derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more
30 sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and

flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories, e.g., for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compounds of general Formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

5 For administration to non-human animals, the composition may also be added to the animal feed or drinking water. It will be convenient to formulate these animal feed and drinking water compositions so that the animal takes in an appropriate quantity of the composition along with its diet. It will also be convenient to present the composition as a premix for addition to the feed or drinking water.

10 Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per human patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain
15 between from about 1 mg to about 500 mg of an active ingredient.

Frequency of dosage may also vary depending on the compound used and the particular disease treated. However, for treatment of most disorders, a dosage regimen of 4 times daily or less is preferred. For the treatment of eating disorders, including obesity, a dosage regimen of 1 or 2 times daily is particularly preferred. For the treatment of impotence
20 a single dose that rapidly reaches effective concentrations is desirable.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular
25 disease undergoing therapy.

Preferred compounds of the invention will have desirable pharmacological properties. Such properties include, but are not limited to oral bioavailability, low toxicity, low serum protein binding and desirable *in vitro* and *in vivo* half-lives. Penetration of the blood brain barrier for compounds used to treat CNS disorders is necessary, while low brain levels of
30 compounds used to treat peripheral disorders are often preferred.

Assays may be used to predict these desirable pharmacological properties. Assays used to predict bioavailability include transport across human intestinal cell monolayers, including Caco-2 cell monolayers. Toxicity to cultured hepatocytes may be used to predict compound toxicity. Penetration of the blood brain barrier of a compound in humans may be

predicted from the brain levels of the compound in laboratory animals given the compound intravenously.

Serum protein binding may be predicted from albumin binding assays. Such assays are described in a review by Oravcová, et al. (Journal of Chromatography B (1996) volume 677, pages 1-27).

Compound half-life is inversely proportional to the frequency of dosage of a compound. *In vitro* half-lives of compounds may be predicted from assays of microsomal half-life as described by Kuhnz and Gieschen (Drug Metabolism and Disposition, (1998) volume 26, pages 1120-1127).

Compounds of Formula I exhibit good activity in standard *in vitro* MCH receptor binding assays and/ or calcium mobilization assays, specifically in the assays as specified in Examples 5 and 6, which follow. References herein to "standard *in vitro* receptor binding assay" are intended to refer to that protocol as defined in Example 5 which follows. References herein to "standard MCH 1 receptor calcium mobilization assay" are intended to refer to that protocol as defined in Example 6 which follows. Generally, preferred compounds of Formula I have an K_i of about 1 micromolar or less, still more preferably a K_i of about 100 nanomolar or less even more preferably a K_i of about 10 nanomolar or less or even 1 nanomolar or less in such a defined standard *in vitro* MCH 1 receptor binding assay and exemplified by Example 5. Generally preferred compounds of Formula I are MCH 1 receptor antagonists and exhibit EC_{50} values of about 4 micromolar or less, more preferably 1 micromolar or less, still more preferably EC_{50} values of about 100 nanomolar or less even more preferably an EC_{50} value of about 10 nanomolar or less or even 1 nanomolar or less in such a defined standard *in vitro* MCH 1 receptor mediated calcium mobilization assay as exemplified by Example 6 which follows.


Preferred compounds of Formula I do not interact with dopamine receptors, particularly human dopamine D2 and D4 receptors. Dopamine receptor binding assays may be preformed using the methods described in Example 9 which follows. Preferred compounds of Formula I exhibit K_i values greater than 1 micromolar in standard assays of dopamine receptor binding assays such as the dopamine D2 and D4 receptor binding assays described in Example 9.

EXAMPLES**Preparation of compounds**


The compounds of the invention can be prepared essentially according to the synthetic procedure shown in Scheme 1. As shown, a 2-phenylacylcycloalkyl compound of general structure 7 may be condensed with a 4-arylpiperazine or piperidine of general structure 8 in the presence of a reducing agent to provide a compound of general Formula I. The reducing agent may be sodium borohydride, sodium triacetoxy borohydride, lithium aluminum hydride, alane or the like. Alternatively, an acid chloride or an acid can be coupled with the piperazine to generate an amide, which can in turn be reduced to yield the desired compound of Formula I.

The preparation of a specific compound of this invention (the 1S,2S enantiomer of Compound 1) is described graphically in Scheme 2 and the synthetic steps used are presented within Example 1. Within Scheme 2, 1S,2S 2-phenylcyclopropanecarboxylic acid (9) was condensed with 1-(4-bromo-3-methoxyphenyl)piperazine (10) in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDCI), dimethylaminopyridine (DMAP) and 1-hydroxybenzotriazole (HOBT). The resulting amide 11 was reduced to the desired amine by reduction with alane in tetrahydrofuran.

Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the invention, as demonstrated by the following examples. Unless otherwise specified all reagents and solvent are of standard commercial grade and are used without further purification.

Scheme 1

Scheme 2

Preparation of 1*S*,2*S* Enantiomer of Compound I

5

Example 1.**Preparation of (1*S*, 2*S*)-1-(4-bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl)methylpiperazine.**

10 Compound numbers 9-11 in the following example represent compounds shown in Scheme 2.

1-(4-Bromo-3-methoxyphenyl)piperazine (10)

A solution of 1-(3-methoxyphenyl)piperazine dihydrobromide (3.5 g, 10 mmol) is dissolved in DMSO (30 mL) and heated at 65 °C for 4h in a flask which is open to the atmosphere. After cooling, the mixture is poured into a separatory funnel containing 100 mL

15

of 1 N sodium hydroxide solution and extracted with ethyl ether (3 X 100 mL). The organic extracts are dried (Na₂SO₄), filtered, and concentrated to provide 1-(4-bromo-3-methoxyphenyl)piperazine as a solid. ¹H NMR (400 MHz, CDCl₃) 7.34-7.36 (d, J=2.2 Hz, 1H), 6.47 (s, 1H), 6.38-6.41 (d, J = 2.8 Hz, 1H), 3.87 (s, 3H, OMe), 3.11-3.13 (m, 4H), 3.01-3.03 (m, 4H)

(1S,2S)-1-(4-bromo-3-methoxyphenyl)-4-(trans-2-phenylcyclopropyl) carbonylpiperazine (11).

EDCI (0.42g, 2.2 mmol), DMAP (0.27 g, 2.2 mmol) and HOBT (0.23 g, 2.2 mmol) are added to a solution of acid 9 (0.34 g, 2.1 mmol) and piperazine 10 (0.54 g, 2.0 mmol) in chloroform (15 mL) and the resulting solution allowed to stir overnight. The solution is washed with water (10 mL), saturated NaHCO₃ solution, (10 mL), brine (10 mL) and dried over magnesium sulfate. After filtration the solution is concentrated and the resulting oil purified by column chromatography eluting with 2% methanol in chloroform to provide the desired amide 11 as a white sticky solid. ¹H NMR (400 MHz, CDCl₃) δ 7.1-7.5 (m, 6H), 6.50 (s, 1H), 6.40 (d, J = 7 Hz, 1H), 3.89 (s, 3H, OMe), 3.8 (bm, 4H), 3.2 (bm, 4H), 2.5 (m, 1H), 1.98 (m, 1H), 1.70 (m, 1H), 1.35 (m, 1H), LCMS (CI) 416 (M+1).

(1S, 2S)-1-(4-bromo-3-methoxyphenyl)-4-(trans-2- phenylcyclopropyl) methylpiperazine (Compound 1, Table 1)

A solution of alane triethylamine complex (3.13 mL, 1.57 mmol) is added to a solution of amide 11 (0.65 g, 1.57 mmol) in THF (10 mL) at 0 °C. After 45 min, the reaction is quenched with water and extracted with ether. The organic extracts are dried (MgSO₄), filtered, and concentrated to a colorless oil which is purified by column chromatography eluting with 5% MeOH/chloroform to provide the desired compound 1 as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.38 (d, J = 7 Hz, 1H), 7.1-7.35 (m, 5H), 6.45 (d, J = 1 Hz, 1H), 6.40 dd, J = 7, 1 Hz, 1H), 3.88 (s, 3H, OMe), 3.2 (m, 4H), 2.7 (m, 4H), 2.63 (m, 1H), 2.40 (dd, J = 12, 5 Hz, 1H), 1.70 (m, 1H), 1.25 (m, 1H), 0.85-1.0 (m, 2H).

Example 2

The following compounds are prepared essentially according to the procedures described with respect to Schemes 1 and 2 and further set forth in Example 1. Variations

suitable for preparing the following compounds will be readily apparent to those skilled in the art of organic synthesis:

- a) 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine
- 5 b) 1*R*, 2*R*-1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine
- c) 1-(4-Iodo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine
- d) 1-(4-Chloro-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine
- e) 1-(4-Phenyl-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine
- f) 1-(4-Bromo-3-methoxyphenyl)-4-[*trans*-2-(3-methoxyphenyl)cyclopropyl]
- 10 methylpiperazine
- g) 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-[4-chlorophenyl] cyclopropyl) methylpiperazine (compound 4)
- h) 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-[2-methylphenyl] cyclopropyl)methylpiperazine (compound 2)
- 15 i) 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-[4-methoxyphenyl] cyclopropyl)methylpiperazine
- j) 1-(4-Bromo-3-methoxyphenyl)-4-([3-phenyl]propen-2-yl)piperazine (Compound 5)
- k) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2-methylphenyl}]propen-2-yl)piperazine (compound 6)
- 20 l) 1-(3-Methoxyphenyl)-4-([3-phenyl]propen-2-yl)piperazine
- m) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3-methylphenyl}]propen-2-yl)piperazine
- n) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2-methoxyphenyl}]propen-2-yl)piperazine
- o) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3-chlorophenyl}]propen-2-yl)piperazine
- p) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3-ethoxyphenyl}]propen-2-yl)piperazine
- 25 q) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2,3-dimethoxyphenyl}]propen-2-yl)piperazine
- r) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3,4-dimethoxyphenyl}]propen-2-yl)piperazine
- s) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2,5-dimethoxyphenyl}]propen-2-yl)piperazine
- t) 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2,4-dimethoxyphenyl}]propen-2-yl)piperazine
- u) 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine
- 30 v) 1-(4-Iodo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine
- w) 1-(4-Chloro-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine
- x) 1-(4-Methyl-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine
- y) 1-(4-Trifluormethyl-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine
- z) 1-(4-Bromo-3-ethoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine.

Example 3.**Cells expressing MCH 1 Receptors**

Cells or preparations of cells recombinantly expressing human MCH 1 receptors,
5 monkey MCH 1 receptors, or chimeric human MCH 1/human Beta 2 Adrenergic receptors
may be used in the radioligand binding assay and Calcium Mobilization assay which follows.
The preparation of expression vectors for such MCH 1 Receptors has been described
previously, e.g. in U.S. Provisional Application No. 60/216,081, filed July 6, 2000 and U.S.
Provisional Application 60/284,835, filed April 19, 2001, pages 19-20 and the sequence
10 listing, both of which application are hereby incorporated by reference for their teachings
regarding the cloning and expression of MCH 1 receptors.

Preparation of HEK 293 cells expressing the monkey MCH receptor

HEK 293 cells are stably transfected via standard calcium phosphate precipitation
procedures with a Cynomolgus macaque monkey MCH expression vector described
15 previously or other MCH 1 receptor expression vector.

Cells are grown to confluency at 37 C, 5% CO₂, approximately 48-72 hours, in
DMEM high glucose culture medium (catalog #10-017-CV, MEDiatech, Herndon, VA)
supplemented with 10% fetal bovine serum, 25 mM HEPES, and 500 ug/ml G418. The cells
are pelleted by gentle centrifugation. Cell pellets are washed twice with cold PBS, harvested
20 in cold PBS containing 5 mM EDTA, and stored at -80 C.

Preparation of CHO cells expressing the monkey MCH receptor

CHO (Chinese Hamster Ovary) cells are transfected via standard calcium phosphate
precipitation procedures with an MCH 1 receptor expression vector.

Cells are grown to confluency at 37 C, 5% CO₂, approximately 48-72 hours, in Ham's
25 F12 culture medium (catalog #10-080-CV, MEDiatech, Herndon, VA) supplemented with
10% fetal bovine serum, 25 mM HEPES, and 500 ug/ml (active) G418. The cells are pelleted
by gentle centrifugation. Cell pellets are washed twice with cold PBS, harvested in cold PBS
containing 5 mM EDTA, and stored at -80 °C.

Example 4.**Purified Membranes**

HEK 293 cell pellets stored frozen at -80 °C are thawed by addition of wash buffer
(25 mM Hepes with 1.0mM CaCl₂, 5.0mM MgCl₂, 120mM NaCl, PH7.4) and homogenized
for 30 seconds using a BRINKMAN POLYTRON, setting 5. Cells are centrifuged for 10

minutes at 48,000 x g. The supernatant is discarded and the pellet is resuspended in fresh wash buffer, and homogenized again. The protein concentration of the resulting membrane preparation is measured using the Bradford protein assay (Bio-Rad Laboratories, Hercules, CA). By this measure, a 1-liter culture of cells typically yields 50-75 mg of total membrane protein.

Example 5.

Radioligand Binding Assays for Modulators of Chimeric Receptors

Purified membranes from HEK 293 cells expressing the monkey MCH receptor are prepared by the procedure given in Example 3. The membrane homogenate is centrifuged as before and resuspended to a protein concentration of 333ug/ml in binding buffer (Wash buffer + 0.1% BSA and 1.0uM final conc. phosphoramidon) for an assay volume of 50ug membrane protein/150ul binding buffer. Phosphoramidon is from SIGMA BIOCHEMICALS, St. Louis, MO (cat# R-7385).

Competition binding assays are performed at room temperature in Falcon 96 well round bottom polypropylene plates. To each assay well is added: 150 ul of MCH receptor containing membranes in binding buffer, prepared as described above, 50 ul ¹²⁵I-Tyr MCH in binding buffer, 50 ul binding buffer, and 2 ul test compound in DMSO. ¹²⁵I-Tyr MCH (specific activity = 2200 Ci/mMol) is purchased from NEN, Boston, MA (Cat # NEX 373) and is diluted in binding buffer to provide a final assay concentration of 30 pM.

Non-specific binding is defined as the binding measured in the presence of 1 uM unlabeled MCH. MCH is purchased from BACHEM U.S.A., King of Prussia, PA (cat # H-1482). To each assay well used to determine non-specific MCH binding is added: 150 ul of MCH receptor-containing membranes in binding buffer, 50 ul ¹²⁵I-Tyr MCH in binding buffer, unlabeled MCH in 25 ul binding buffer, and 25 ul binding buffer.

Assay plates are incubated for 1 hour at room temperature. Membranes are harvested onto WALLAC glass fiber filters (PERKIN-ELMER, Gaithersburg, MD) which are pre-soaked with 1.0% PEI (polyethyleneimine) for 2 hours prior to use. Filters are allowed to dry overnight then counted in a WALLAC 1205 BETA PLATE counter after addition of WALLAC BETA SCINT scintillation fluid.

For saturation binding the concentration of ¹²⁵I-Tyr MCH is varied from 7 – 1,000 pM. Typically 11 concentration points are collected per saturation binding curve.

Equilibrium binding parameters are determined by fitting the allosteric Hill equation to the measured values with the aid of the computer program FitP™ (BIOSOFT, Ferguson, MO).

5 **Example 6.**

Functional Assay of Monkey MCH receptors

Calcium mobilization assay

The following assay can be used to monitor the response of cells expressing melanin concentrating hormone receptors to melanin concentrating hormone. The assay can also be
10 used to determine if test compounds act as agonists or antagonists of melanin concentrating hormone receptors.

Chinese Hamster Ovary (CHO) cells stably transfected with an MCH 1 receptor expression vector are grown to a density of 15,000 cells/well in FALCON black-walled, clear-bottomed 96-well plates (#3904, BECTON-DICKINSON, Franklin Lakes, NJ). Prior to
15 running the assay the culture medium is emptied from the 96 well plates. Fluo-3 calcium sensitive dye (Molecular Probes, Eugene, OR) is added to each well (dye solution: 1 mg FLUO-3 AM, 440 uL DMSO and 440 ul 20% pluronic acid in DMSO, diluted 1:4, 50 ul diluted solution per well). Plates are covered with aluminum foil and incubated at 37°C for 1-2 hours. After the incubation the dye solution is emptied from the plates, cells are washed
20 once in 100 ul KRH buffer (0.05 mM KCl, 0.115 M NaCl, 9.6 mM NaH₂PO₄, 0.01 mM MgSO₄, 25 mM HEPES, pH 7.4) to remove excess dye; after washing 80 ul KRH buffer is added to each well.

Determination of Agonist Effects

Fluorescence response may monitored upon the addition of either human MCH or test
25 compound as described below by a FLIPR™ plate reader (Molecular Devices, Sunnyvale, CA) by excitation at 480 nM and emission at 530 nM.

Determination of Antagonist Effects

In order to measure the ability of a test compound to antagonize the response of cells expressing MCH receptors to MCH, the EC₅₀ of MCH is first determined.

30 An additional 20 ul of KRH buffer and 1 ul DMSO is added to each well of cells, prepared as described immediately above. 100 ul human MCH in KRH buffer is automatically transferred by the FLIPR instrument to each well. An 8-point concentration response curve, with final MCH concentrations of 1 nM to 3 uM, is used to determine MCH EC₅₀.

Test compounds are dissolved in DMSO, diluted in 20 ul KRH buffer, and added to

cells prepared as described above. The 96 well plates containing prepared cells and test compounds are incubated in the dark, at room temperature for 0.5 – 6 hours. It is important that the incubation not continue beyond 6 hours. Just prior to determining the fluorescence response 100 ul human MCH diluted in KRH buffer to 2 x EC₅₀ is automatically added by the

5 FLIPR instrument to each well of the 96 well plate for a final sample volume of 200 ul and a final MCH concentration of EC₅₀. The final concentration of test compounds in the assay wells is between 1 uM and 5 uM. Typically cells exposed to one EC₅₀ of MCH exhibit a fluorescence response of about 10,000 Relative Fluorescence Units. Antagonists of the MCH receptor exhibit a response that is significantly less than that of the control cells to the p≤0.05

10 level, as measured using a parametric test of statistical significance. Typically antagonists of the MCH receptor decrease the fluorescence response relative to control cells by about 20%, preferably by about 50%, and most preferably by at least 80% as compared to matched controls.

Determination of Agonist Effects

15 The ability of a compound to act as an agonist of the MCH receptor may be determined by measuring the fluorescence response of cells expressing MCH receptors, using the methods described above, in the absence of MCH. Compounds that cause cells to exhibit fluorescence above background are MCH 1 receptor agonists.

20 Example 7

Preparation of radiolabeled probe compounds of the invention

The compounds of the invention are prepared as radiolabeled probes by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope. The radioisotope is preferably selected from of at least one of carbon (preferably ¹⁴C), hydrogen

25 (preferably ³H), sulfur (preferably ³⁵S), or iodine (preferably ¹²⁵I). Such radiolabeled probes are conveniently synthesized by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds. Such suppliers include Amersham Corporation, Arlington Heights, IL; Cambridge Isotope Laboratories, Inc. Andover, MA; SRI International, Menlo Park, CA; Wizard Laboratories, West Sacramento, CA; ChemSyn Laboratories, Lexena, KS;

30 American Radiolabeled Chemicals, Inc., St. Louis, MO; and Moravsek Biochemicals Inc., Brea, CA.

Tritium labeled probe compounds are also conveniently prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas. Such preparations

are also conveniently carried out as a custom radiolabeling by any of the suppliers listed in the preceding paragraph using the compound of the invention as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as appropriate.

5

Example 8

Use of compounds of the invention as probes for melanin receptors in cultured cells and tissue samples

Receptor autoradiography (receptor mapping) of melanin concentrating hormone
10 receptors in cultured cells or tissue samples is carried out in vitro as described by Kuhar in sections 8.1.1 to 8.1.9 of Current Protocols in Pharmacology (1998) John Wiley & Sons, New York, using radiolabeled compounds of the invention prepared as described in the preceding Example.

15 Example 9

Determination of D₂ and D₄ receptor binding activity

The following assay is a standard assay for determining the binding affinity of compounds to dopamine D₄ and D₂ receptors.

Pellets of Chinese hamster ovary (CHO) cells containing recombinantly expressing
20 primate D₂, human D₄ dopamine receptors are used for the assays. The sample is homogenized in 100 volumes (w/vol) of 0.05 M Tris HCl buffer containing 120 mM NaCl, 5 mM MgCl₂ and 1 mM EDTA at 4°C and pH 7.4. The sample is then centrifuged at 30,000 x g and resuspended and rehomogenized. The sample is then centrifuged as described and the final tissue sample is frozen until use. The tissue is resuspended 1:20 (wt/vol) in 0.05 M Tris
25 HCl buffer containing 120 mM NaCl.

Incubations for dopaminergic binding are carried out at 25°C and contain 0.4 ml of tissue sample, 0.1 nM ³H-YM 09151-2 (Nemonapride, cis-5-Chloro-2-methoxy-4-(methylamino)-N-(2-methyl-2-(phenylmethyl)-3-pyrrolidinyl)benzamide) and the compound of interest in a total incubation of 1.0 ml. Nonspecific binding is defined as that binding
30 found in the presence of 1 micromolar spiperone; without further additions, nonspecific binding is less than 20% of total binding.


The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which

it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the invention and that modifications may be made therein without departing from the spirit or scope of the invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims

5 conclude this specification.


What is claimed is:

1. A compound of the formula:



or a pharmaceutically acceptable salt thereof wherein:

- 5 Q represents a group of the Formula:



wherein A is C₁-C₅ alkylene optionally mono-, di, or trisubstituted with substituents independently chosen from C₁-C₃ alkyl, C₁-C₃ alkoxy, halogen, halo(C₁-C₃)alkyl, halo(C₁-C₃)alkoxy, hydroxy, amino, and mono- or di(C₁-C₃)alkylamino;

- 10 R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are the same or different and represent hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -SO₂NH₂, mono or dialkylsulfonamido, -C(O)NH₂, or mono or di(C₁-C₆)alkylcarboxamido;
- 15 R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ independently represent hydrogen or C₁-C₆ alkyl;

W is nitrogen or C-R_a where R_a represents hydrogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl or cyano;

X represents halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkylamino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di(C₁-C₆)alkylcarboxamido, -SO₂NH₂, mono or di(C₁-C₆)alkylsulfonamido; or

X represents phenyl which may be optionally substituted with up to five substituents, which are the same or different and are selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-


C₆alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxy carbonyl, -COOH, -CONH₂, mono- or di-(C₁-C₆)alkylcarboxamido, -SO₂NH₂, and mono or di(C₁-C₆)alkylsulfonamido;

Y is oxygen, sulfur, -S(O)-, or -SO₂-; and

Z is C₁-C₆ alkyl or mono, di or trifluoromethyl.

5

2. A compound or salt according to claim 1, wherein Q is a group of the Formula:



and A is methylene optionally substituted with C₁-C₂ alkyl.

10

3. A compound or salt according to claim 2, wherein A is methylene and W is nitrogen.

4. A compound or salt according to claim 3, wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen.

15

5. A compound or salt according to claim 4, wherein:
wherein

X is halogen;

Y is oxygen; and

20

Z is C₁-C₆ alkyl.

6. A compound or salt according to claim 4, wherein:
R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;

25

X is hydrogen, halogen, or phenyl;

Y is oxygen; and

Z is C₁-C₆ alkyl.

30

7. A compound or salt according to claim 4, wherein:
R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;

X is halogen;

Y is oxygen; and

Z is C₁-C₆ alkyl.

8. A compound or salt according to claim 2, wherein A is methylene and W is
5 CH.

9. A compound or salt according to claim 8, wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅,
R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen.

10. A compound or salt according to claim 9, wherein:

X is halogen;

Y is oxygen; and

Z is C₁-C₆ alkyl.

11. A compound or salt according to claim 9, wherein:

R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen,
halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;

X is hydrogen, halogen, or phenyl;

Y is oxygen; and

20 Z is C₁-C₆ alkyl.

12. A compound or salt according to claim 9, wherein:


R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen,
halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;

25 X is halogen;

Y is oxygen; and

Z is C₁-C₆ alkyl.

13. A compound or salt according to Claim 2, of the formula



where A is methylene optionally substituted with C₁-C₂ alkyl.


14. A compound or salt according to Claim 13, wherein A is methylene and W is
5 nitrogen or CH.

15. A compound or salt according to claim 14, wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₅, R₁₇,
R₁₈, and R₁₉ are hydrogen.


10 16. A compound or salt according to claim 15, wherein
R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ independently represent hydrogen, halogen, C₁-C₆ alkyl, C₁-
C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;
R₁₄ and R₁₆ are the same or different and are either hydrogen or methyl;
X is hydrogen, halogen, or phenyl;
15 Y is oxygen; and
Z is C₁-C₆ alkyl.

17. A compound or salt according to Claim 14, wherein
R₁, R₆, R₇, R₈, R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ are hydrogen;
20 R₂, R₃, R₄, and R₅ are independently hydrogen, C₁-C₂ alkyl, C₁-C₂ alkoxy, or halogen;
X is halogen;
Y is oxygen; and
Z is C₁-C₆ alkyl.

25 18. A compound or salt according to claim 1 of the formula



19. A compound or salt according to claim 18, wherein:
- 5 R₁₃, R₁₅, R₁₇, R₁₉, are hydrogen; and
 R₁₀, R₁₁, R₁₂, R₁₄, R₁₆, and R₁₈ independently represent hydrogen or methyl.
20. A compound or salt according to claim 19, wherein R₁₀, R₁₁, R₁₂, R₁₄, R₁₆, and
 10 R₁₈ are hydrogen.
21. A compound or salt according to claim 20, wherein W is N or CH.
22. A compound or salt according to claim 21, wherein:
- R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ independently represent hydrogen, halogen, C₁-C₆ alkyl,
 15 C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;
 X is hydrogen or halogen;
 Y is oxygen; and
 Z is C₁-C₆ alkyl.
- 20 23. A compound or salt according to claim 21, wherein:
- R₁, R₂, R₃, R₄ independently represent hydrogen, halogen, C₁-C₂alkyl, or C₁-C₂ alkoxy;
 R₅, R₆, R₇, and R₈ are hydrogen;
 X is halogen;
 Y is oxygen; and
 25 Z is C₁-C₆ alkyl.
24. A compound or salt according to claim 1, of the formula



25. A compound or salt according to claim 24, wherein:
R₁₃, R₁₅, R₁₇, R₁₉, are hydrogen; and
5 R₁₀, R₁₁, R₁₂, R₁₄, R₁₆, and R₁₈ independently represent hydrogen or methyl.
26. A compound or salt according to claim 25, wherein R₁₀, R₁₁, R₁₂, R₁₄, R₁₆, and R₁₈ are hydrogen.
- 10 27. A compound or salt according to claim 26, wherein W is N or CH.
28. A compound or salt according to claim 27 wherein
R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ independently represent hydrogen, halogen, C₁-C₆ alkyl,
C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;
15 X is hydrogen or halogen;
Y is oxygen; and
Z is C₁-C₆ alkyl.
29. A compound or salt according to Claim 27 wherein
20 R₁, R₂, R₃, R₄ independently represent hydrogen, halogen, C₁-C₂alkyl, or C₁-C₂ alkoxy;
R₅, R₆, R₇, and R₈ are hydrogen;
X is halogen;
Y is oxygen; and
Z is C₁-C₆ alkyl.
- 25 30. A compound or salt according to Claim 1, which is selected from:
1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine;
(1*S*, 2*S*)-1-(4-bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl)methylpiperazine;


- 1*R*, 2*R*-1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine;
 1-(4-Iodo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine;
 1-(4-Chloro-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine;
 1-(4-Phenyl-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperazine;
 5 1-(4-Bromo-3-methoxyphenyl)-4-[*trans*-2-(3-methoxyphenyl)cyclopropyl] methylpiperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-[4-chlorophenyl] cyclopropyl) methylpiperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-[2-methylphenyl] cyclopropyl)methylpiperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-[4-methoxyphenyl] cyclopropyl)methylpiperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine;
 10 1-(4-Iodo-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine;
 1-(4-Chloro-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine;
 1-(4-Methyl-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine;
 1-(4-Trifluormethyl-3-methoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine; and
 1-(4-Bromo-3-ethoxyphenyl)-4-(*trans*-2-phenylcyclopropyl) methylpiperidine; or a
 15 pharmaceutically acceptable salt thereof.

31. A compound or salt according to Claim 1, which is selected from:

- 1-(4-Bromo-3-methoxyphenyl)-4-([3-phenyl]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2-methylphenyl}]propen-2-yl)piperazine;
 20 1-(3-Methoxyphenyl)-4-([3-phenyl]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3-methylphenyl}]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2-methoxyphenyl}]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3-chlorophenyl}]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3-ethoxyphenyl}]propen-2-yl)piperazine;
 25 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2,3-dimethoxyphenyl}]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{3,4-dimethoxyphenyl}]propen-2-yl)piperazine;
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2,5-dimethoxyphenyl}]propen-2-yl)piperazine; and
 1-(4-Bromo-3-methoxyphenyl)-4-([3-{2,4-dimethoxyphenyl}]propen-2-yl)piperazine, or a
 pharmaceutically acceptable salt thereof.


30

32. A compound of the formula:



wherein:

Q represents a group of the Formula:




- 5 wherein A is C₁-C₅ alkylene optionally mono-, di, or trisubstituted with substituents independently chosen from C₁-C₃ alkyl, C₁-C₃ alkoxy, halogen, halo(C₁-C₃)alkyl, halo(C₁-C₃)alkoxy, hydroxy, amino, and mono- or di(C₁-C₃)alkylamino;
- R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ are the same or different and represent hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -SO₂NH₂, mono or dialkylsulfonamido, -C(O)NH₂, or mono or di(C₁-C₆)alkylcarboxamido;
- 10 R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, R₁₈, and R₁₉ independently represent hydrogen or C₁-C₆ alkyl;
- 15 W is nitrogen or C-R_a where R_a represents hydrogen, hydroxy, C₁-C₆ alkoxy, C₁-C₆ alkyl or cyano;
- X represents halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkylamino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di(C₁-C₆)alkylcarboxamido, -SO₂NH₂, mono or di(C₁-C₆)alkylsulfonamido; or
- 20 X represents phenyl which may be optionally substituted by up to five substituents, which may be the same or different and are selected from the group consisting of hydrogen, halogen, cyano, nitro, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ alkoxy, C₁-C₆

alkylthio, hydroxy, amino, mono or di(C₁-C₆)alkyl amino, halo(C₁-C₆)alkyl, halo(C₁-C₆)alkoxy,

C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, -COOH, -CONH₂, mono- or di-(C₁-C₆)alkylcarboxamido, -SO₂NH₂, and mono or di(C₁-C₆)alkylsulfonamido;

- 5 Y is oxygen, sulfur, -S(O)-, or -SO₂-; and
Z is C₁-C₆ alkyl or mono, di or trifluoromethyl.

33. A compound or salt according to claim 32, wherein Q is a group the formula



- 10 where A is methylene optionally substituted with C₁-C₂ alkyl or A is a single bond.

34. A compound according to claim 32, wherein W is nitrogen or CH.

35. A compound according to claim 34, wherein R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇,
15 R₁₈, and R₁₉ are hydrogen.

36. A compound according to claim 35 wherein:

X is halogen;

Y is oxygen; and

- 20 Z is C₁-C₆ alkyl.

37. A compound according to Claim 35 wherein

R₁, R₂, R₃, R₄, R₅, R₆, R₇, and R₈ may be the same or different and represent hydrogen, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or trifluoromethoxy;

- 25 X is halogen;
Y is oxygen; and
Z is C₁-C₆ alkyl.

38. A pharmaceutical composition comprising a compound or salt according Claim 1
30 combined with at least one pharmaceutically acceptable carrier or excipient.

39. The pharmaceutical composition of Claim 38 wherein the pharmaceutical composition is formulated as an injectable fluid, a pill, a capsule, a syrup, or a transdermal patch.

5 40. A method for the treatment of obesity, said method comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of Claim 1.

41. A method for demonstrating the presence of MCH 1 receptors in cell or tissue
10 samples, said method comprising:

preparing a plurality of matched cell or tissue samples,

preparing at least one control sample by contacting under conditions that permit
binding of MCH to MCH 1 receptors within cell and tissue samples at least one of the
matched cell or tissue samples with a control solution comprising a detectably-labeled
15 preparation of a selected compound or salt of Claims 1 at a first measured molar
concentration, said control solution further comprising an unlabelled preparation of the
selected compound or salt at a second measured molar concentration, which second measured
concentration is greater than said first measured concentration,

preparing at least one experimental sample by contacting under conditions that permit
20 binding of MCH to MCH 1 receptors within cell and tissue samples at least one of the
matched cell or tissue samples with an experimental solution comprising the detectably-
labeled preparation of the selected compound or salt at the first measured molar
concentration, said experimental solution not further comprising an unlabelled preparation of
any compound or salt of any of Claim 1 at a concentration greater than or equal to said first
25 measured concentration;

washing the at least one control sample to remove unbound selected compound or salt
to produce at least one washed control sample;

washing the at least one experimental sample to remove unbound selected compound
or salt to produce at least one washed experimental sample;

30 measuring the amount of detectable label of any remaining bound detectably-labeled
selected compound or salt in the at least one washed control sample;

measuring the amount detectable label of any remaining bound detectably-labeled
selected compound or salt in the at least one washed experimental sample;

comparing the amount of detectable label measured in each of the at least one washed experimental sample to the amount of detectable label measured in each of the at least one washed control sample

wherein, a comparison that indicates the detection of a greater amount of detectable label in the at least one washed experimental sample than is detected in any of the at least one washed control samples demonstrates the presence of MCH 1 receptors in that experimental sample.

42. The method of Claim 41 wherein the compound is radiolabeled.

43. The method of Claim 42 wherein the detection is accomplished using autoradiography.

44. A method for altering the signal-transducing activity of MCH 1 receptors, said method comprising contacting cells expressing such receptors with a solution comprising a compound according to Claim 1 at a concentration sufficient to detectably alter the electrophysiology of the cell, wherein a detectable alteration of the electrophysiology of the cell indicates an alteration of the signal-transducing activity of MCH 1 receptors.

45. The method of Claim 44 wherein the cell is a neuronal cell that is contacted in vivo in an animal, the solution is a body fluid, and the alteration in the electrophysiology of the cell is detected as a reproducible change in the animal's feeding behavior.

46. The method of Claim 45 wherein the animal is a human, the cell is a brain cell, and the fluid is cerebrospinal fluid.

47. A packaged pharmaceutical composition comprising the pharmaceutical composition of Claim 38 in a container and instructions for using the composition to treat a patient suffering from obesity.

48. The use of a compound according to any one of claims 1-37 in the preparation of a medicament for use in the treatment of obesity.

THIS PAGE BLANK (USPTO)